# ESPC Coupled Global Prediction System Develop and Test Coupled Physical Parameterizations: NAVGEM/CICE/HYCOM

Pamela Posey Naval Research Laboratory (NRL) Stennis Space Center, MS

phone: (228) 688-5596 fax: (228) 688-4759 email: pamela.posey@nrlmry.navy.mil

Shouping Wang Naval Research Laboratory (NRL) Monterey, CA

phone: (831) 656-4719 fax: (831) 656-4769 email: wang@nrlmry.navy.mil

Award Number: N0001413WX20376 http://www7320.nrlssc.navy.mil/hycomARC

Team consists of: James Chen, Alan Wallcraft, Joe Metzger, Pam Posey and Shouping Wang

## **LONG-TERM GOALS**

A fully coupled global atmosphere/wave/ocean/land/ice prediction system providing daily predications out to 10 days and weekly predictions out to 30 days.

# **OBJECTIVES**

The initial operational capability of a fully coupled global atmosphere/wave/ocean/land/ice prediction system is targeted for 2018. Predictions will provide environmental information to meet Navy and DoD operations and planning needs throughout the globe from undersea to the upper atmosphere and from the tropics to the poles. The system will be implemented on Navy operational computer systems, and the necessary processing infrastructure will be put in place to provide products for the Navy fleet user consumption.

### **APPROACH**

Building on the ESMF NUOPC interoperability layers, present advanced models representing the physics within each of the earth components of atmosphere, ocean, surface waves, ice and land will pass information through separate flux computation models. The system will use analysis fields of each component as initial conditions and make daily forecasts out to 10 days. Throughout each weekly cycle, predictions out to 30 days will be constructed.

Task 1). Implementation of coupling physics between NAVGEM/HYCOM/CICE.

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send commentarters Services, Directorate for Inf	ts regarding this burden estimate formation Operations and Reports	or any other aspect of the property of the pro	his collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 30 SEP 2013	2 DEDORT TVD			3. DATES COVERED <b>00-00-2013 to 00-00-2013</b>	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
ESPC Coupled Global Prediction System - Develop and Test Coupled Physical Parameterizations: NAVGEM/CICE/HYCOM				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Naval Research Laboratory, Stennis Space Center, MS, 39529				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release; distributi	ion unlimited			
13. SUPPLEMENTARY NO	OTES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF	
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	7	RESPONSIBLE PERSON

**Report Documentation Page** 

Form Approved OMB No. 0704-0188

## WORK COMPLETED

At the start of this project, HYCOM and CICE have already been coupled using the Earth System Modeling Framework (ESMF) with atmospheric forcing input from files and the results validated. Summer seasonal forecasts (out to 5 months) have been run as part of the Study of the Environmental Arctic Change (SEARCH) Sea Ice Outlook (SIO) effort. The SIO is an international effort to provide a community-wide summary of the expected September arctic sea ice minimum. Monthly reports released throughout the summer synthesize community estimates of the current state and expected minimum of sea ice.

Along with the backbone components of this system (NAVGEM/HYCOM/CICE), other data models have been used to read archived atmospheric data, ocean data and other observed data. Figure 1 shows a system of CICE coupled with Data\_Atmos and Data\_Ocean. Figure 2 shows a system of CICE coupled with Data\_Atmos, Data\_Ocean and Data\_Assim (assimilation).




Figure 1. A system of CICE coupled with Data Atmos and Data Ocean.




Figure 2. A system of CICE coupled with Data\_Atmos, Data\_Ocean and Data\_Assim (assimilation).

## **RESULTS**

In May 2013, the Arctic Cap Nowcast/Forecast System (ACNFS), i.e. coupled HYCOM/CICE, was run in forward model mode, without assimilation, initialized with a May 1, 2013 analysis to determine a seasonal projection of September Arctic sea ice minimum. Eight model runs were made, using Navy atmospheric forcing from 2003-2012 and run forward for 5 months for each specific year. This ensemble of eight members gives an indication of how sea ice can respond to variable atmospheric conditions during summer. Ice extent was calculated using all grid cells with at least 15% ice concentration and then averaged over each day in September. Ice extent averaged across all enesemble members during September was the September ice extent extimate. For the runs initialzed from May 01, 2013, the ACNFS outlook sea ice extent estimate was 4.9 Mkm² +/- 0.7 Mkm². Figure 3 shows the distribution of the individual Pan-Arctic Outlook values (June report) for September 2013 sea ice extent. The 2013 sea ice minimum was 4.81 Mkm² observed on September 13.




Figure 3. Distribution of individual Pan-Arctic Outlook values (June Report) for September 2013 sea ice extent (values are rounded to the tenths).

We have also tested a single column model of the arctic surface flux parameterization. It uses the mosaic method to calculate the fluxes in summer conditions where sea ice and melting ponds may coexist. Tests show that the heat and moisture roughness are sensitive to the sea surface conditions. Figure 4 shows that both the momentum and scalar roughness lengths are strong functions of wind speed over sea ice. The ratios of the scalars to the momentum roughness are also dependent on the surface dynamic condition. Conventionally, the roughness lengths are specified as constants in the numerical models such as in GFS and COAMPS; this relationship should be correctly modified in both the ESPC efforts and within the operational NAVGEM system according to the new understanding.




Figure 4. Representation of surface roughness over sea ice following Andreas et al. (2010). Left: roughness of momentum  $(z_0)$ , heat  $(z_{0h})$  and moisture  $(z_{0q})$  as a function of wind speed; right: the ratio  $z_{0h}/z_0$  or  $z_{0q}/z_0$  as a function of the bulk roughness Reynolds number.

An example of a coupled run product from the Data\_Atmos-Data\_Ocean-CICE system (Figure 1) is shown in Figure 5, in which contours of sea ice concentration (SIC) and sea surface temperture (SST) at the end of 1440 hrs (60 days) are displayed. To validate with the observed data, we use GHRSST analyzed SIC for comparison as shown in Figure 6. In general, at the end of the two month spin-up simulation , CICE can generate a result which is in good agreement with the observed GHRSST SIC data.




Figure 5. 1440 hour (60 day, from 01 Jan 2010) sea ice concentration (SIC) simulated by the Data\_Atmos-Data\_Ocean-CICE system.




Figure 6. GHRSST Sea Ice Concentration (SIC) analysis. The display time is the same time as that in Figure 5.

Using the four-model system (Data\_Atmos, Data\_Ocean, Data\_Assim and CICE – shown in Figure 2), a spinup for sea ice using NOGAPS and HYCOM archived historic data are employed to drive CICE. Moreover, we use GHRSST analyzed sea ice concentration (SIC) data to assimilate sea ice into CICE via Data\_Assim model. The assimilated result is shown in Figure 7.




Figure 7. Same as in Figure 5, except the run assimilated GHRSST analyzed sea ice concentration.

Note, the corresponding non-assimilated case is shown in Figure 5. To compare with the GHRSST observed sea ice concentration data (Figure 6), we found that the assimilated case has done its job which is better than that of the non-assimilated case.

The plan is to use the CICE restart file generated by the ice spin-up procedure (using the Data\_Atmos-Data\_Ocean-CICE system) as the initial condition for the NAVGEM-Data\_Ocean-CICE or NAVGEM-HYCOM-CICE systems to conduct forecasts.

### **IMPACT/APPLICATIONS**

Surface heat fluxes and winds must be exchanged between NAVGEM, HYCOM and CICE. The fluxes and winds must be implemented for ESPC through the NUOPC interoperability layer. Consideration of flux computations from both the atmosphere and the ice perspectives must be taken into account and move toward consistent flux computations. Additional effects will also be added, including variations in albedo and roughness on wind drag.

### RELATED PROJECTS

The five other subprojects in ESPC project:

- 1) Coupling infrastructure and interoperability layer extension across all earth components
- 2) Develop and test coupled physical parameterizations: NAVGEM/HYCOM
- 3) Develop and test coupled physical parameterizations and tropolar wave model grid: NAVGEM/ WaveWatch-III/HYCOM
- 4) Aerosol Physical Coupling
- 5) Development of ESMF for coupling capability (NOAA/ESRL)

## REFERENCES

Andreas, E.L., T. W. Horst, A. A. Grachev P. O. Persson, C. W. Fairall, P. S. Guest, and R. E. Jordan, 2010, Parameterizing turbulent exchange over summer sea ice and the marginal ice zone. *Q. J. R. Meteorol. Soc.* 136: 927–943.

# **PUBLICATIONS**

P.G. Posey, E. J. Metzger, R. A. Allard, R. H. Preller, D. Hebert, A.J. Wallcraft, M.W. Phelps, and O. M. Smedstad, "Arctic Cap Nowcast Forecast System (ACNFS) End of Summer 2013 Ice Extent Projection- June Report", Sea Ice Outlook Project website, posted Jun, Jul, Aug and Sept, 2013.